arXiv:math/0311386v2 [math.DS] 13 Aug 2004

THE MARKUS-YAMABE CONJECTURE FOR DIFFERENTIABLE VECTOR FIELDS OF $\mathbb{R}^2$.

CARLOS GUTIERREZ AND ROLAND RABANAL

Dedicated to César Camacho on his 60th Birthday

Abstract. (a) Let $X\colon \mathbb{R}^2 \to \mathbb{R}^2$ be a differentiable map (not necessarily C^1) and let $\mathrm{Spec}(X)$ be the set of (complex) eigenvalues of the derivative DX_p when p varies in $\mathbb{R}^2$. If, for some $\epsilon > 0$, $\mathrm{Spec}(X) \cap [0, \epsilon) = \emptyset$ then X is injective.

(b) Let $X\colon \mathbb{R}^2 \to \mathbb{R}^2$ be a differentiable vector field such that $X(0) = 0$ and $\mathrm{Spec}(X) \subset \{z \in \mathbb{C} : \Re(z) < 0\}$. Then, for all $p \in \mathbb{R}^2$, there is a unique positive trajectory starting at p; moreover the $\omega-$limit set of p is equal to $\{0\}$.

Carlos Gutierrez and Roland Rabanal
Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo, caixa postal 668, 13560-970, São Carlos SP, Brasil
E-mail address: gutp@icmc.usp.br, roland@icmc.usp.br